

Volunteer Lake Assessment Program Individual Lake Reports MASSASECUM, LAKE, BRADFORD, NH

MORPHOMETRIC DATA							<u>CLASSIFICATION</u>	KNOWN EXOTIC SPECIES	
Watershed Area (Ac.):	6,044	Max. Depth (m):	16.4	Flushing Rate (yr¹)	2	Year	Trophic class	Variable Milfoil	
Surface Area (Ac.):	402	Mean Depth (m):	3.8	P Retention Coef:	0.59	1987	MESOTROPHIC		
Shore Length (m):	6,400	Volume (m³):	6,229,000	Elevation (ft):	631	2005	MESOTROPHIC		

The Waterbody Report Card tables are generated from the 2012 305(b) report on the status of N.H. waters, and are based on data collected from 2001-2011.

Designated Use	Parameter	Category	Comments				
Aquatic Life	Phosphorus (Total)	Very Good	>5 samples and median is < 1/2 threshold.				
	рН	Slightly Bad	>10% of samples exceed criteria by a small margin (minimum of 2 exceedances).				
	D.O. (mg/L)	Encouraging	< 10 samples and no exceedance of criteria. More data needed.				
	D.O. (% sat)	Slightly Bad	>10% of samples exceed criteria by a small margin (minimum of 2 exceedances).				
	Chlorophyll-a	Good	>/=5 samples and median is < threshold but > 1/2 threshold value.				
Primary Contact Recreation	E. coli	Very Good	All bacteria samples <75% of geometric mean criteria, but not enough to calculate geometric mean. Or, all bacteria samples are < single sample criteria and calculated Geometric means are less than geometric mean criteria.				
	Chlorophyll-a	Very Good	At least 10 samples with 0 exceedances of criteria.				

BEACH PRIMARY CONTACT ASSESSMENT STATUS

LAKE MASSASECUM - CAMP PIESAULE BEACH	E. coli	Very Good	All bacteria samples <75% of geometric mean criteria, but not enough to calculate geometric mean. Or, all bacteria samples are < single sample criteria and calculated Geometric means are less than geometric mean criteria.
LAKE MASSASECUM - FRENCH'S PARK TOWN BEACH	E. coli	Slightly Bad	Slightly exceeds criteria.
LAKE MASSASECUM - MASSASECUM CASINO BEACH	E. coli	Very Good	All bacteria samples <75% of geometric mean criteria, but not enough to calculate geometric mean. Or, all bacteria samples are < single sample criteria and calculated Geometric means are less than geometric mean criteria.

WATERSHED LAND USE SUMMARY

Fry, J., Xian, G., Jin, S., Dewitz, J., Homer, C., Yang, L., Barnes, C., Herold, N., and Wickham, J., 2011. Completion of the 2006 National Land Cover Database for the Conterminous United States, PERS, Vol. 77(9):858-864. For larger image contact NHDES.



Land Cover Category % Cover		Land Cover Category	% Cover	Land Cover Category	% Cover	
Open Water	7.14	Barren Land	0	Grassland/Herbaceous	1.22	
Developed-Open Space	3.22	Deciduous Forest	27.57	Pasture Hay	0.81	
Developed-Low Intensity	0.67	Evergreen Forest	28.23	Cultivated Crops	0	
Developed-Medium Intensity	0	Mixed Forest	28.68	Woody Wetlands	1.96	
Developed-High Intensity	0	Shrub-Scrub	0.43	Emergent Wetlands	0.13	



VOLUNTEER LAKE ASSESSMENT PROGRAM INDIVIDUAL LAKE REPORTS LAKE MASSASECUM, BRADFORD, NH 2012 DATA SUMMARY

OBSERVATIONS AND RECOMMENDATIONS (Refer to Table 1 and Historical Deep Spot Data Graphic)

- **CHLOROPHYLL-A:** Chlorophyll levels were relatively low and less than the NH lake median. Historical trend analysis indicates a relatively stable chlorophyll level since monitoring began.
- **CONDUCTIVITY/CHLORIDE:** Conductivity levels were elevated in Frenches Park and Davis Brooks.
- **E. COLI:** E. coli levels were well below the state standard for surface waters.
- Total Phosphorus: Deep spot phosphorus levels were low and historical trend analysis indicates a relatively stable epilimnetic (upper water layer) phosphorus level since monitoring began. Davis Brook phosphorus levels were slightly elevated likely due to wetland influences. Howlett Brook phosphorus was slightly elevated in August likely due to sediments as turbidity was also elevated.
- Transparency: Transparency improved in August and average transparency was greater than the NH lake median. Historical trend analysis indicates a relatively stable transparency since monitoring began.
- TURBIDITY: Turbidity was elevated in Howlett Brook in August likely due to low stream flows and sediment contamination.
- PH: Deep spot pH levels were lower than desirable and potentially critical to aquatic life.
- RECOMMENDED ACTIONS: Conduct chloride monitoring at Davis and Frenches Park Brooks to establish baseline data sets. Keep up the great work!

	Table 1. 2012 Average Water Quality Data for MASSASECUM LAKE										
	Alk.	Chlor-a ug/l	Chloride mg/l	Cond. uS/cm	E. Coli #/100ml	Total P ug/l	Trans. m		Turb. ntu	pН	
Station Name	mg/l										
							NVS	VS			
Babcook Brook				50.4		12			0.45	6.57	
Colby Brook				30.3		14			0.43	6.83	
Davis Brook				89.7	70	27			1.46	6.09	
Davis Brook #2					25						
Deep Epilimnion	3.9	3.87	5	39.9		5	5.25	5.63	0.48	6.42	
Deep Metalimnion				40.1		8			0.61	5.94	
Deep Hypolimnion				43.2		9			1.83	5.97	
Frenches Park Brook				168.8		6			0.84	6.70	
Howlett Brook				42.9		17			1.95	6.87	
Melvin Bk Outlet				41.5		7			0.59	6.74	

NH Median Values: Median values for specific parameters generated from historic lake monitoring

data.

Alkalinity: 4.9 mg/L Chlorophyll-a: 4.58 mg/m³ Conductivity: 40.0 uS/cm Chloride: 4 mg/L Total Phosphorus: 12 ug/L Transparency: 3.2 m

pH: 6.6

NH Water Quality Standards: Numeric criteria for specific parameters. Results exceeding criteria are considered a

water quality violation.

Chloride: < 230 mg/L (chronic)

E. coli: > 88 cts/100 mL – public beach
E. coli: > 406 cts/100 mL – surface waters
Turbidity: > 10 NTU above natural level
pH: 6.5-8.0 (unless naturally occurring)

HISTORICAL WATER QUALITY TREND ANALYSIS

ParameterTrendExplanationChlorophyll-aStableData not significantly increasing or decreasing.TransparencyStableData not significantly increasing or decreasing.Phosphorus (epilimnion)StableData not significantly increasing or decreasing.

This report was generated by the NH DES Volunteer Lake Assessment Program (VLAP). For more information contact: Sara Steiner

PO Box 95 Concord, NH 03302-0095 (603) 271-2658 sara.steiner@des.nh.gov



